

Recombinant Human EXOSC10 293 Cell Lysate

Cat. No. EXOSC10-6504HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for exosome component 10 (EXOSC10), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

EXOSC10 exosome component 10 [Homo sapiens]

Official Symbol

EXOSC10

Synonyms

EXOSC10; exosome component 10; PMSCL2, polymyositis/scleroderma autoantigen 2, 100kDa; p2; p3; p4; PM Scl; PM/Scl 100; polymyositis/scleroderma autoantigen 2 (100kD); RRP6; Rrp6p; autoantigen PM-SCL; autoantigen PM/Scl 2; polymyositis/scleroderma autoantigen 100 kDa; polymyositis/scleroderma autoantigen 2, 100kDa; P100 polymyositis-scleroderma overlap syndrome-associated autoantigen; PMSCL; PM-Scl; PMSCL2; PM/Scl-100;

Gene ID

5394

mRNA Refseq

NM_001001998

Protein Refseq

NP_001001998

MIM

605960

UniProt ID

Q01780

Chromosome Location

1p36.22

Pathway

RNA degradation, organism-specific biosystem; RNA degradation, conserved

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem;

Function

3-5 exonuclease activity; RNA binding; exoribonuclease activity; hydrolase activity; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA