

Recombinant Human ESD 293 Cell Lysate

Cat. No. ESD-6541HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for esterase D/formylglutathione hydrolase (ESD) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name ESD esterase D [Homo sapiens]

Official Symbol ESD

Synonyms ESD; esterase D; esterase D/formylglutathione hydrolase; S-formylglutathione hydrolase; S formylglutathione hydrolase; esterase 10; methylumbelliferyl-acetate deacetylase; FGH; FLJ11763;

Gene ID 2098

mRNA Refseq NM_001984

Protein Refseq NP_001975

MIM 133280

UniProt ID P10768

Chromosome Location 13q14.1-q14.2

Pathway formaldehyde oxidation II (glutathione-dependent), organism-specific biosystem;

Function S-formylglutathione hydrolase activity; carboxylesterase activity; hydrolase activity, acting on ester bonds; methylumbelliferyl-acetate deacetylase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA