

Recombinant Human EPHX1 293 Cell Lysate

Cat. No. EPHX1-6586HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for epoxide hydrolase 1, microsomal (xenobiotic) (EPHX1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	EPHX1 epoxide hydrolase 1, microsomal (xenobiotic) [Homo sapiens]
Official Symbol	EPHX1
Synonyms	EPHX1; epoxide hydrolase 1, microsomal (xenobiotic); EPHX; epoxide hydrolase 1; epoxide hydratase; MEH; EPOX; HYL1;
Gene ID	2052
mRNA Refseq	NM_000120
Protein Refseq	NP_000111
UniProt ID	P07099
Chromosome Location	1q42.1
Pathway	Bile secretion, organism-specific biosystem; Bile secretion, conserved biosystem; Metabolism of xenobiotics by cytochrome P450, organism-specific biosystem; Metabolism of xenobiotics by cytochrome P450, conserved biosystem;
Function	cis-stilbene-oxide hydrolase activity; epoxide hydrolase activity; hydrolase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA