

Recombinant Human EIF5A 293 Cell Lysate

Cat. No. EIF5A-6641HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 5A (EIF5A), transcript variant B is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	EIF5A eukaryotic translation initiation factor 5A [Homo sapiens]
Official Symbol	EIF5A
Synonyms	EIF5A; eukaryotic translation initiation factor 5A; eukaryotic translation initiation factor 5A-1; EIF 5A; EIF5A1; MGC99547; MGC104255; eIF-4D; eIF-5A1; eIF-5A-1; rev-binding factor; eukaryotic initiation factor 5A; EIF-5A; eIF5A1;
Gene ID	1984
mRNA Refseq	NM_001143760
Protein Refseq	NP_001137232
MIM	600187
UniProt ID	P63241
Chromosome Location	17p13-p12
Pathway	Hypusine synthesis from eIF5A-lysine, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; PTM: gamma carboxylation, hypusine formation and arylsulfatase activation, organism-specific biosystem; Post-translational protein modification, organism-specific biosystem; Translation Factors, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem;

Function

RNA binding; U6 snRNA binding; protein N-terminus binding; protein binding;
ribosome binding; translation elongation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA