

Recombinant Human EIF4A1 293 Cell Lysate

Cat. No. EIF4A1-6654HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 4A1 (EIF4A1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

EIF4A1 eukaryotic translation initiation factor 4A1 [Homo sapiens]

Official Symbol

EIF4A1

Synonyms

EIF4A1; eukaryotic translation initiation factor 4A1; EIF4A, eukaryotic translation initiation factor 4A, isoform 1; eukaryotic initiation factor 4A-I; DDX2A; EIF 4A; eukaryotic initiation factor 4AI; ATP-dependent RNA helicase eIF4A-1; eukaryotic translation initiation factor 4A, isoform 1; EIF4A; EIF-4A; eIF4A-I; eIF-4A-I;

Gene ID

1973

mRNA Refseq

NM_001204510

Protein Refseq

NP_001191439

MIM

602641

UniProt ID

P60842

Chromosome Location

17p13

Pathway

Activation of the mRNA upon binding of the cap-binding complex and eIFs, and subsequent binding to 43S, organism-specific biosystem; Antiviral mechanism by IFN-stimulated genes, organism-specific biosystem; Cap-dependent Translation Initiation,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; Cytokine Signaling in Immune system, organism-specific biosystem; Deadenylation of mRNA, organism-specific biosystem; Deadenylation-dependent mRNA decay, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem;

Function

ATP binding; ATP-dependent helicase activity; RNA cap binding; helicase activity; hydrolase activity; mRNA binding; nucleotide binding; protein binding; translation factor activity, nucleic acid binding; translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA