

Recombinant Human EIF3M 293 Cell Lysate

Cat. No. EIF3M-6655HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 3, subunit M (EIF3M) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name EIF3M eukaryotic translation initiation factor 3, subunit M [Homo sapiens]

Official Symbol EIF3M

Synonyms EIF3M; eukaryotic translation initiation factor 3, subunit M; PCI domain containing 1 (herpesvirus entry mediator) , PCID1; eukaryotic translation initiation factor 3 subunit M; eIF3m; FLJ29030; GA17; hfl B5; B5 receptor; fetal lung protein B5; dendritic cell protein; PCI domain-containing protein 1; PCI domain containing 1 (herpesvirus entry mediator); B5; PCID1; hfl-B5;

Gene ID 10480

mRNA Refseq NM_006360

Protein Refseq NP_006351

MIM 609641

UniProt ID Q7L2H7

Chromosome Location 11p13

Function protein binding; contributes_to translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA