

Recombinant Human EIF3J 293 Cell Lysate

Cat. No. EIF3J-6657HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for eukaryotic translation initiation factor 3, subunit J (EIF3J) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

EIF3J eukaryotic translation initiation factor 3, subunit J [Homo sapiens]

Official Symbol

EIF3J

Synonyms

EIF3J; eukaryotic translation initiation factor 3, subunit J; EIF3S1, eukaryotic translation initiation factor 3, subunit 1 alpha, 35kDa; eukaryotic translation initiation factor 3 subunit J; eIF3 alpha; eIF3 p35; eIF3j; eIF-3-alpha; eukaryotic translation initiation factor 3 subunit 1; eukaryotic translation initiation factor 3, subunit 1 alpha, 35kDa; eukaryotic translation initiation factor 3, subunit 1 (alpha, 35kD); EIF3S1; eIF3-p35; eIF3-alpha;

Gene ID

8669

mRNA Refseq

NM_003758

Protein Refseq

NP_003749

MIM

603910

UniProt ID

O75822

Chromosome Location

15q21.1

Pathway

Activation of the mRNA upon binding of the cap-binding complex and eIFs, and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

subsequent binding to 43S, organism-specific biosystem; Cap-dependent Translation Initiation, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem; Formation of a pool of free 40S subunits, organism-specific biosystem; Formation of the ternary complex, and subsequently, the 43S complex, organism-specific biosystem; GTP hydrolysis and joining of the 60S ribosomal subunit, organism-specific biosystem; Gene Expression, organism-specific biosystem;

Function

protein binding; contributes_to translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA