

Recombinant Human EIF3H 293 Cell Lysate

Cat. No. EIF3H-6659HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 3, subunit H (EIF3H) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

EIF3H eukaryotic translation initiation factor 3, subunit H [Homo sapiens]

Official Symbol

EIF3H

Synonyms

EIF3H; eukaryotic translation initiation factor 3, subunit H; EIF3S3, eukaryotic translation initiation factor 3, subunit 3 gamma, 40kDa; eukaryotic translation initiation factor 3 subunit H; eIF3 gamma; eIF3 p40; eIF3h; eIF-3-gamma; eIF3 p40 subunit; eukaryotic translation initiation factor 3 subunit 3; eukaryotic translation initiation factor 3, subunit 2 (beta, 36kD); eukaryotic translation initiation factor 3, subunit 3 gamma, 40kDa; eukaryotic translation initiation factor 3, subunit 3 (gamma, 40kD); EIF3S3; eIF3-p40; eIF3-gamma; MGC102958;

Gene ID

8667

mRNA Refseq

NM_003756

Protein Refseq

NP_003747

MIM

603912

UniProt ID

O15372

**Chromosome
Location**

8q24.11

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Activation of the mRNA upon binding of the cap-binding complex and eIFs, and subsequent binding to 43S, organism-specific biosystem; Cap-dependent Translation Initiation, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem; Formation of a pool of free 40S subunits, organism-specific biosystem; Formation of the ternary complex, and subsequently, the 43S complex, organism-specific biosystem; GTP hydrolysis and joining of the 60S ribosomal subunit, organism-specific biosystem; Gene Expression, organism-specific biosystem;

Function

protein binding; contributes_to translation initiation factor activity; contributes_to translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA