

Recombinant Human EIF3F 293 Cell Lysate

Cat. No. EIF3F-6661HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 3, subunit F (EIF3F) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name EIF3F eukaryotic translation initiation factor 3, subunit F [Homo sapiens]

Official Symbol EIF3F

Synonyms EIF3F; eukaryotic translation initiation factor 3, subunit F; EIF3S5, eukaryotic translation initiation factor 3, subunit 5 epsilon, 47kDa; eukaryotic translation initiation factor 3 subunit F; eIF3 epsilon; eIF3 p47; eIF3f; eIF3-epsilon; eIF-3-epsilon; eukaryotic translation initiation factor 3, subunit 5 epsilon, 47kDa; eukaryotic translation initiation factor 3, subunit 5 (epsilon, 47kD); EIF3S5; eIF3-p47;

Gene ID 8665

mRNA Refseq NM_003754

Protein Refseq NP_003745

MIM 603914

UniProt ID O00303

Chromosome Location 11p15.4

Pathway Activation of the mRNA upon binding of the cap-binding complex and eIFs, and subsequent binding to 43S, organism-specific biosystem; Cap-dependent Translation

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Initiation, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem; Formation of a pool of free 40S subunits, organism-specific biosystem; Formation of the ternary complex, and subsequently, the 43S complex, organism-specific biosystem; GTP hydrolysis and joining of the 60S ribosomal subunit, organism-specific biosystem; Gene Expression, organism-specific biosystem;

Function

protein binding; contributes_to translation initiation factor activity; contributes_to translation initiation factor activity; translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA