

Recombinant Human DLG3 293 Cell Lysate

Cat. No. DLG3-6911HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for discs, large homolog 3 (Drosophila) (DLG3), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

DLG3 discs, large homolog 3 (Drosophila) [Homo sapiens]

Official Symbol

DLG3

Synonyms

DLG3; discs, large homolog 3 (Drosophila); discs, large homolog 3 (neuroendocrine dlG, Drosophila); disks large homolog 3; KIAA1232; MRX90; NE DIg; NEDLG; neuroendocrine dlG; SAP 102; SAP102; neuroendocrine-DLG; synapse-associated protein 102; MRX; XLMR;

Gene ID

1741

mRNA Refseq

NM_001166278

Protein Refseq

NP_001159750

MIM

300189

UniProt ID

Q92796

Chromosome Location

Xq13.1

Pathway

Activation of Ca-permeable Kainate Receptor, organism-specific biosystem; Activation of Kainate Receptors upon glutamate binding, organism-specific biosystem; Axon guidance, organism-specific biosystem; Developmental Biology,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

organism-specific biosystem; Ionotropic activity of Kainate Receptors, organism-specific biosystem; L1CAM interactions, organism-specific biosystem; Neuronal System, organism-specific biosystem;

Function

PDZ domain binding; guanylate kinase activity; ionotropic glutamate receptor binding; phosphatase binding; protein C-terminus binding; protein binding; protein phosphatase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA