

Recombinant Human CYP27A1 293 Cell Lysate

Cat. No. CYP27A1-7119HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for cytochrome P450, family 27, subfamily A, polypeptide 1 (CYP27A1), nuclear gene encoding mitochondrial protein is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CYP27A1 cytochrome P450, family 27, subfamily A, polypeptide 1 [Homo sapiens]

Official Symbol

CYP27A1

Synonyms

CYP27A1; cytochrome P450, family 27, subfamily A, polypeptide 1; CYP27, cytochrome P450, subfamily XXVIIA (steroid 27 hydroxylase, cerebrotendinous xanthomatosis), polypeptide 1; sterol 26-hydroxylase, mitochondrial; cerebrotendinous xanthomatosis; CP27; CTX; cytochrome P450 27; sterol 27-hydroxylase; cytochrome P-450C27/25; vitamin D(3) 25-hydroxylase; cholestanetriol 26-monooxygenase; 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol 27-hydroxylase; 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha-triol 26-hydroxylase; 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha-triol 27-hydroxylase; cytochrome P450, subfamily XXVIIA (steroid 27-hydroxylase, cerebrotendinous xanthomatosis), polypeptide 1; CYP27;

Gene ID

1593

mRNA Refseq

NM_000784

Protein Refseq

NP_000775

MIM

606530

UniProt ID

Q02318

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	2q33-qter
Pathway	Bile acid and bile salt metabolism, organism-specific biosystem; Bile acid biosynthesis, cholesterol =>cholate, organism-specific biosystem; Bile acid biosynthesis, cholesterol => cholate, conserved biosystem; Biological oxidations, organism-specific biosystem; Cytochrome P450 - arranged by substrate type, organism-specific biosystem;
Function	cholestanetriol 26-monooxygenase activity; cholesterol 26-hydroxylase activity; electron carrier activity; heme binding; metal ion binding; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen; steroid hydroxylase activity; vitamin D3 25-hydroxylase activity;