

Recombinant Human CYP26A1 293 Cell Lysate

Cat. No. CYP26A1-7121HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for cytochrome P450, family 26, subfamily A, polypeptide 1 (CYP26A1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [CYP26A1 cytochrome P450, family 26, subfamily A, polypeptide 1 \[Homo sapiens \]](#)

Official Symbol CYP26A1

Synonyms CYP26A1; cytochrome P450, family 26, subfamily A, polypeptide 1; cytochrome P450, subfamily XXVIA, polypeptide 1; cytochrome P450 26A1; CP26; CYP26; P450RAI; P450RAI1; hP450RAI; cytochrome P450RAI; retinoic acid 4-hydroxylase; P450, retinoic acid-inactivating, 1; retinoic acid-metabolizing cytochrome; cytochrome P450 retinoic acid-inactivating 1;

Gene ID [1592](#)

mRNA Refseq [NM_000783](#)

Protein Refseq [NP_000774](#)

MIM [602239](#)

UniProt ID [O43174](#)

Chromosome Location 10q23-q24

Pathway Adipogenesis, organism-specific biosystem; Biological oxidations, organism-specific biosystem; Cytochrome P450 - arranged by substrate type, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Metabolism, organism-specific biosystem; Nuclear receptors in lipid metabolism and toxicity, organism-specific biosystem; Phase 1 - Functionalization of compounds, organism-specific biosystem; Retinol metabolism, organism-specific biosystem;

Function

electron carrier activity; heme binding; heme binding; metal ion binding; monooxygenase activity; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen; oxygen binding; retinoic acid 4-hydroxylase activity; retinoic acid binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA