

Recombinant Human CYP19A1 293 Cell Lysate

Cat. No. CYP19A1-7127HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for cytochrome P450, family 19, subfamily A, polypeptide 1 (CYP19A1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CYP19A1 cytochrome P450, family 19, subfamily A, polypeptide 1 [Homo sapiens]

Official Symbol

CYP19A1

Synonyms

CYP19A1; cytochrome P450, family 19, subfamily A, polypeptide 1; CYP19, cytochrome P450, subfamily XIX (aromatization of androgens); cytochrome P450 19A1; ARO; ARO1; aromatase; CPV1; CYAR; P 450AROM; estrogen synthase; estrogen synthetase; cytochrome P-450AROM; microsomal monooxygenase; flavoprotein-linked monooxygenase; cytochrome P450, subfamily XIX (aromatization of androgens); CYP19; CYPXIX; P-450AROM; MGC104309;

Gene ID

1588

mRNA Refseq

NM_000103

Protein Refseq

NP_000094

MIM

107910

UniProt ID

P11511

Chromosome Location

15q21

Pathway

Biological oxidations, organism-specific biosystem; C19/C18-Steroid hormone

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosynthesis, pregnenolone => androstenedione => estrone, organism-specific biosystem; C19/C18-Steroid hormone biosynthesis, pregnenolone => androstenedione => estrone, conserved biosystem;

Function

aromatase activity; electron carrier activity; electron carrier activity; heme binding; heme binding; metal ion binding; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen, reduced flavin or flavoprotein as one donor, and incorporation of one atom of oxygen; oxygen binding;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA