

Recombinant Human CYP17A1 293 Cell Lysate

Cat. No. CYP17A1-7128HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for cytochrome P450, family 17, subfamily A, polypeptide 1 (CYP17A1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CYP17A1 cytochrome P450, family 17, subfamily A, polypeptide 1 [Homo sapiens]

Official Symbol

CYP17A1

Synonyms

CYP17A1; cytochrome P450, family 17, subfamily A, polypeptide 1; CYP17, cytochrome P450, subfamily XVII (steroid 17 alpha hydroxylase), adrenal hyperplasia; steroid 17-alpha-hydroxylase/17,20 lyase; CPT7; P450C17; S17AH; Steroid 17 alpha monooxygenase; CYPXVII; cytochrome P450c17; cytochrome P450-C17; cytochrome P450 17A1; cytochrome p450 XVIIA1; steroid 17-alpha-monooxygenase; cytochrome P450, subfamily XVII (steroid 17-alpha-hydroxylase), adrenal hyperplasia; CYP17;

Gene ID

1586

mRNA Refseq

NM_000102

Protein Refseq

NP_000093

MIM

609300

UniProt ID

P05093

**Chromosome
Location**

10q24.3

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Androgen biosynthesis, organism-specific biosystem; Biological oxidations, organism-specific biosystem; C19/C18-Steroid hormone biosynthesis, pregnenolone => androstenedione => estrone, organism-specific biosystem; C19/C18-Steroid hormone biosynthesis, pregnenolone => androstenedione =>
Function	electron carrier activity; heme binding; metal ion binding; monooxygenase activity; oxygen binding; steroid 17-alpha-monooxygenase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA