Recombinant Human CXCR4 293 Cell Lysate

Cat. No. CXCR4-7161HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-X-C motif) receptor 4 (CXCR4), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CXCR4 chemokine (C-X-C motif) receptor 4 [Homo sapiens]
Official Symbol	CXCR4
Synonyms	CXCR4; chemokine (C-X-C motif) receptor 4; chemokine (C X C motif), receptor 4 (fusin); C-X-C chemokine receptor type 4; CD184; D2S201E; fusin; HM89; HSY3RR; LESTR; NPY3R; NPYR; NPYY3R; CXC-R4; CXCR-4; CD184 antigen; SDF-1 receptor; neuropeptide Y receptor Y3; seven transmembrane helix receptor; stromal cell-derived factor 1 receptor; lipopolysaccharide-associated protein 3; seven-transmembrane-segment receptor, spleen; leukocyte-derived seven transmembrane domain receptor; leukocyte-derived seven-transmembrane-domain receptor; FB22; LAP3; LCR1; WHIM; NPYRL;
Gene ID	7852
mRNA Refseq	NM_001008540
Protein Refseq	NP_001008540
MIM	162643
UniProt ID	P61073
Chromosome Location	2q21

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem; Binding and entry of HIV virion, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem;
Function	C-X-C chemokine receptor activity; G-protein coupled receptor activity; actin binding; coreceptor activity; myosin light chain binding; protein binding; receptor activity; signal transducer activity; ubiquitin binding; ubiquitin protein ligase binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA