

Recombinant Human CXCL9 293 Cell Lysate

Cat. No. CXCL9-7165HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-X-C motif) ligand 9 (CXCL9) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CXCL9 chemokine (C-X-C motif) ligand 9 [Homo sapiens]

Official Symbol

CXCL9

Synonyms

CXCL9; chemokine (C-X-C motif) ligand 9; CMK, MIG, monokine induced by gamma interferon; C-X-C motif chemokine 9; crg 10; Humig; SCYB9; small-inducible cytokine B9; gamma-interferon-induced monokine; monokine induced by gamma interferon; monokine induced by interferon-gamma; CMK; MIG; crg-10;

Gene ID

4283

mRNA Refseq

NM_002416

Protein Refseq

NP_002407

MIM

601704

UniProt ID

Q07325

Chromosome Location

4q21

Pathway

CXCR3-mediated signaling events, organism-specific biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem;
Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-
cytokine receptor interaction, conserved biosystem;

Function

chemokine activity; cytokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA