

Recombinant Human CWC27 293 Cell Lysate

Cat. No. CWC27-7174HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for CWC27 spliceosome-associated protein homolog (<i>S. cerevisiae</i>) (CWC27) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CWC27 CWC27 spliceosome-associated protein homolog (S. cerevisiae) [Homo sapiens]

Official Symbol

CWC27

Synonyms

CWC27; CWC27 spliceosome-associated protein homolog (S. cerevisiae); SDCCAG10, serologically defined colon cancer antigen 10; peptidyl-prolyl cis-trans isomerase CWC27 homolog; NY CO 10; PPIase CWC27; PPIase SDCCAG10; antigen NY-CO-10; peptidyl-prolyl cis-trans isomerase SDCCAG10; serologically defined colon cancer antigen 10; NY-CO-10; SDCCAG10;

Gene ID

10283

mRNA Refseq

NM_005869

Protein Refseq

NP_005860

UniProt ID

Q6UX04

Chromosome Location

5q12.3

Function

isomerase activity; peptidyl-prolyl cis-trans isomerase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA