

Recombinant Human CUL3 293 Cell Lysate

Cat. No. CUL3-7183HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for cullin 3 (CUL3) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CUL3 cullin 3 [Homo sapiens]
Official Symbol	CUL3
Synonyms	CUL3; cullin 3; cullin-3; CUL-3; PHA2E; FLJ25665;
Gene ID	8452
mRNA Refseq	NM_001257197
Protein Refseq	NP_001244126
MIM	603136
UniProt ID	Q13618
Chromosome Location	2q36.2
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and Proteasome degradation, organism-specific biosystem; Aurora B signaling, organism-specific biosystem; Canonical Wnt signaling pathway, organism-specific biosystem; Class I MHC mediated antigen processing &
Function	cyclin binding; protein binding; ubiquitin protein ligase binding; contributes_to

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ubiquitin-protein ligase activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA