Recombinant Human CREB1 293 Cell Lysate

Cat. No. CREB1-7289HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for cAMP responsive element binding protein 1 (CREB1), transcript variant A is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CREB1 cAMP responsive element binding protein 1 [Homo sapiens]
Official Symbol	CREB1
Synonyms	CREB1; cAMP responsive element binding protein 1; cyclic AMP-responsive element-binding protein 1; CREB-1; transactivator protein; active transcription factor CREB; cAMP-response element-binding protein-1; cAMP-responsive element-binding protein 1; CREB; MGC9284;
Gene ID	1385
mRNA Refseq	NM_004379
Protein Refseq	NP_004370
MIM	123810
UniProt ID	P16220
Chromosome Location	2q34
Pathway	AKT phosphorylates targets in the nucleus, organism-specific biosystem; ATF-2 transcription factor network, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Activation of NMDA receptor upon glutamate binding

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	and postsynaptic events, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Adipogenesis, organism-specific biosystem; Amphetamine addiction, organism-specific biosystem;
Function	RNA polymerase II activating transcription factor binding; RNA polymerase II distal enhancer sequence-specific DNA binding; RNA polymerase II transcription factor binding transcription factor activity involved in positive regulation of transcription; cAMP response element binding; double-stranded DNA binding; protein binding; protein dimerization activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; sequence-specific distal enhancer binding RNA polymerase II transcription factor activity; transcription cofactor activity; transcription factor binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA