

Recombinant Human CEP63 293 Cell Lysate

Cat. No. CEP63-7571HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for centrosomal protein 63kDa (CEP63), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [CEP63 centrosomal protein 63kDa \[Homo sapiens \]](#)

Official Symbol CEP63

Synonyms centrosomal protein 63kDa; FLJ13386; centrosomal protein of 63 kDa; centrosome protein CEP63; Cep63

Gene ID [80254](#)

mRNA Refseq [NM_001042383.1](#)

Protein Refseq [NP_001035842.1](#)

MIM [614724](#)

UniProt ID [Q96MT8](#)

Chromosome Location 3q22.2

Pathway Cell Cycle, organism-specific biosystem; Cell Cycle, Mitotic, organism-specific biosystem; Centrosome maturation, organism-specific biosystem; G2/M Transition, organism-specific biosystem; Loss of Nlp from mitotic centrosomes, organism-specific biosystem; Loss of proteins required for interphase microtubule organization??from the centrosome, organism-specific biosystem; Mitotic G2-G2/M phases, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



specific biosystem;

Function

protein binding

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA