

Recombinant Human CDK7 293 Cell Lysate

Cat. No. CDK7-7621HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for cyclin-dependent kinase 7 (CDK7) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CDK7 cyclin-dependent kinase 7 [Homo sapiens]

Official Symbol

CDK7

Synonyms

CDK7; cyclin-dependent kinase 7; cyclin dependent kinase 7 (homolog of Xenopus MO15 cdk activating kinase) , cyclin dependent kinase 7 (MO15 homolog, Xenopus laevis, cdk activating kinase); CAK; CAK1; CDKN7; MO15; STK1; p39 Mo15; protein kinase; 39 KDa protein kinase; kinase subunit of CAK; CDK-activating kinase 1; serine/threonine kinase stk1; cell division protein kinase 7; serine/threonine protein kinase 1; serine/threonine-protein kinase 1; serine/threonine protein kinase MO15; homolog of Xenopus MO15 Cdk-activating kinase; TFIIH basal transcription factor complex kinase subunit; cyclin-dependent kinase 7 (MO15 homolog, Xenopus laevis, cdk-activating kinase); HCAK; p39MO15;

Gene ID

1022

mRNA Refseq

NM_001799

Protein Refseq

NP_001790

MIM

601955

UniProt ID

P50613

Chromosome

5q12.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Androgen Receptor Signaling Pathway, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; Basal transcription factors, organism-specific biosystem; Basal transcription factors, conserved biosystem; Cell Cycle, organism-specific biosystem; Cell Cycle, Mitotic, organism-specific biosystem; Cell cycle, organism-specific biosystem;
Function	ATP binding; DNA-dependent ATPase activity; RNA polymerase II carboxy-terminal domain kinase activity; androgen receptor binding; cyclin-dependent protein kinase activity; nucleotide binding; protein C-terminus binding; protein complex binding; protein kinase activity; transcription coactivator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA