

Recombinant Human CDC27 293 Cell Lysate

Cat. No. CDC27-7662HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for cell division cycle 27 homolog (*S. cerevisiae*) (CDC27), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CDC27 cell division cycle 27 homolog (*S. cerevisiae*) [*Homo sapiens*]

Official Symbol

CDC27

Synonyms

CDC27; cell division cycle 27 homolog (*S. cerevisiae*); cell division cycle 27 , D17S978E, D0S1430E; cell division cycle protein 27 homolog; ANAPC3; anaphase promoting complex subunit 3; APC3; NUC2; H-NUC; nuc2 homolog; CDC27 homolog; anaphase-promoting complex subunit 3; anaphase-promoting complex, protein 3; HNUC; CDC27Hs; D0S1430E; D17S978E;

Gene ID

996

mRNA Refseq

NM_001114091

Protein Refseq

NP_001107563

MIM

116946

UniProt ID

P30260

Chromosome Location

17q21.32

Pathway

APC/C complex, organism-specific biosystem; APC/C complex, conserved biosystem; APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

APC/C:Cdc20 mediated degradation of Cyclin B, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;
APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted
proteins in late mitosis/early G1, organism-specific biosystem;

Function

protein binding; protein phosphatase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA