Recombinant Human CCT8 293 Cell Lysate

Cat. No. CCT8-7686HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for chaperonin containing TCP1, subunit 8 (theta) (CCT8) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CCT8 chaperonin containing TCP1, subunit 8 (theta) [Homo sapiens]
Official Symbol	CCT8
Synonyms	CCT8; chaperonin containing TCP1, subunit 8 (theta); C21orf112, chromosome 21 open reading frame 112; T-complex protein 1 subunit theta; Cctq; PRED71; CCT-theta; TCP-1-theta; renal carcinoma antigen NY-REN-15; T-complex protein 1, theta subunit; D21S246; C21orf112; KIAA0002;
Gene ID	10694
mRNA Refseq	NM_006585
Protein Refseq	NP_006576
UniProt ID	P50990
Chromosome Location	21q21.3
Pathway	Association of TriC/CCT with target proteins during biosynthesis, organism-specific biosystem; Chaperonin-mediated protein folding, organism-specific biosystem; Cooperation of Prefoldin and TriC/CCTin actin and tubulin folding, organism-specific biosystem; Folding of actin by CCT/TriC, organism-specific biosystem; Formation of tubulin folding intermediates by CCT/TriC, organism-specific biosystem; Metabolism

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	of proteins, organism-specific biosystem; Prefoldin mediated transfer of substrateto CCT/TriC, organism-specific biosystem;
Function	ATP binding; ATPase activity, coupled; nucleotide binding; unfolded protein binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA