

Recombinant Human CCT8 293 Cell Lysate

Cat. No. CCT8-7686HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for chaperonin containing TCP1, subunit 8 (theta) (CCT8) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CCT8 chaperonin containing TCP1, subunit 8 (theta) [Homo sapiens]

Official Symbol

CCT8

Synonyms

CCT8; chaperonin containing TCP1, subunit 8 (theta); C21orf112, chromosome 21 open reading frame 112; T-complex protein 1 subunit theta; Cctq; PRED71; CCT-theta; TCP-1-theta; renal carcinoma antigen NY-REN-15; T-complex protein 1, theta subunit; D21S246; C21orf112; KIAA0002;

Gene ID

10694

mRNA Refseq

NM_006585

Protein Refseq

NP_006576

UniProt ID

P50990

Chromosome Location

21q21.3

Pathway

Association of TriC/CCT with target proteins during biosynthesis, organism-specific biosystem; Chaperonin-mediated protein folding, organism-specific biosystem; Cooperation of Prefoldin and TriC/CCTin actin and tubulin folding, organism-specific biosystem; Folding of actin by CCT/TriC, organism-specific biosystem; Formation of tubulin folding intermediates by CCT/TriC, organism-specific biosystem; Metabolism

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

of proteins, organism-specific biosystem; Prefoldin mediated transfer of substrate to CCT/TriC, organism-specific biosystem;

Function

ATP binding; ATPase activity, coupled; nucleotide binding; unfolded protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA