

Recombinant Human CCT3 293 Cell Lysate

Cat. No. CCT3-7690HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for chaperonin containing TCP1, subunit 3 (gamma) (CCT3), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CCT3 chaperonin containing TCP1, subunit 3 (gamma) [Homo sapiens]
Official Symbol	CCT3
Synonyms	CCT3; chaperonin containing TCP1, subunit 3 (gamma); TRiC5; T-complex protein 1 subunit gamma; Cctg; hTRiC5; T-complex protein 1, gamma subunit; TCP1 (t-complex-1) ring complex, polypeptide 5; CCTG; PIG48; CCT-gamma; TCP-1-gamma;
Gene ID	7203
mRNA Refseq	NM_005998
Protein Refseq	NP_005989
MIM	600114
UniProt ID	P49368
Chromosome Location	1q23
Pathway	Association of TriC/CCT with target proteins during biosynthesis, organism-specific biosystem; Chaperonin-mediated protein folding, organism-specific biosystem; Cooperation of Prefoldin and TriC/CCTin actin and tubulin folding, organism-specific biosystem; Folding of actin by CCT/TriC, organism-specific biosystem; Formation of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

tubulin folding intermediates by CCT/TriC, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; Prefoldin mediated transfer of substrate to CCT/TriC, organism-specific biosystem;

Function

ATP binding; nucleotide binding; protein binding; unfolded protein binding;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA