

Recombinant Human CCT2 293 Cell Lysate

Cat. No. CCT2-7691HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for chaperonin containing TCP1, subunit 2 (beta) (CCT2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CCT2 chaperonin containing TCP1, subunit 2 (beta) [Homo sapiens]

Official Symbol

CCT2

Synonyms

CCT2; chaperonin containing TCP1, subunit 2 (beta); T-complex protein 1 subunit beta; Cctb; T-complex protein 1, beta subunit; chaperonin containing t-complex polypeptide 1, subunit 2; chaperonin containing t-complex polypeptide 1, beta subunit; CCTB; 99D8.1; PRO1633; CCT-beta; TCP-1-beta; MGC142074; MGC142076;

Gene ID

10576

mRNA Refseq

NM_001198842

Protein Refseq

NP_001185771

MIM

605139

UniProt ID

P78371

**Chromosome
Location**

12q14

Pathway

Association of TriC/CCT with target proteins during biosynthesis, organism-specific biosystem; Chaperonin-mediated protein folding, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Cooperation of Prefoldin and TriC/CCTin actin and tubulin folding, organism-specific biosystem; Folding of actin by CCT/TriC, organism-specific biosystem; Formation of tubulin folding intermediates by CCT/TriC, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; Prefoldin mediated transfer of substrateto CCT/TriC, organism-specific biosystem;

Function

ATP binding; nucleotide binding; unfolded protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA