Recombinant Human CCL4L1 293 Cell Lysate

Cat. No. CCL4L1-7720HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-C motif) ligand 4-like 1 (CCL4L1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	CCL4L1 chemokine (C-C motif) ligand 4-like 1 [Homo sapiens]
Official Symbol	CCL4L1
Synonyms	CCL4L1; chemokine (C-C motif) ligand 4-like 1; CCL4L, chemokine (C C motif) ligand 4 like , SCYA4L, small inducible cytokine A4 like; C-C motif chemokine 4-like; AT744.2; LAG 1; MIP-1-beta; macrophage inflammatory protein-1b2; lymphocyte activation gene 1 protein; macrophage inflammatory protein 1-beta; monocyte adherence-induced protein 5-alpha; LAG1; CCL4L; LAG-1; SCYA4L;
Gene ID	9560
mRNA Refseq	NM_001001435
Protein Refseq	NP_001001435
MIM	603782
UniProt ID	Q8NHW4
Chromosome Location	17q12
Pathway	Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Cytosolic DNA-sensing pathway, organism-specific biosystem; Cytosolic DNA-sensing pathway, conserved biosystem; IL12-mediated signaling events, organism-specific biosystem;
Function	chemokine activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA