

Recombinant Human CCL26 293 Cell Lysate

Cat. No. CCL26-7725HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-C motif) ligand 26 (CCL26) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name CCL26 chemokine (C-C motif) ligand 26 [Homo sapiens]

Official Symbol CCL26

Synonyms CCL26; chemokine (C-C motif) ligand 26; SCYA26, small inducible cytokine subfamily A (Cys Cys), member 26; C-C motif chemokine 26; CC chemokine IMAC; chemokine N1; eotaxin 3; IMAC; macrophage inflammatory protein 4 alpha; MIP 4a; MIP 4alpha; small inducible cytokine A26; thymic stroma chemokine 1; TSC 1; eotaxin-3; MIP-4-alpha; thymic stroma chemokine-1; small-inducible cytokine A26; macrophage inflammatory protein 4-alpha; small inducible cytokine subfamily A (Cys-Cys), member 26; TSC-1; MIP-4a; SCYA26; MIP-4alpha; MGC126714;

Gene ID 10344

mRNA Refseq NM_006072

Protein Refseq NP_006063

MIM 604697

UniProt ID Q9Y258

Chromosome Location 7q11.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; IL4-mediated signaling events, organism-specific biosystem;
Function	chemokine activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA