

Recombinant Human CCL14 293 Cell Lysate

Cat. No. CCL14-7732HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for chemokine (C-C motif) ligand 14 (CCL14), transcript variant 3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CCL14 chemokine (C-C motif) ligand 14 [Homo sapiens]

Official Symbol

CCL14

Synonyms

CCL14; chemokine (C-C motif) ligand 14; SCYA14, small inducible cytokine subfamily A (Cys Cys), member 14; C-C motif chemokine 14; CKb1; HCC 1; HCC 3; MCIF; NCC 2; SCYL2; chemokine CC-3; new CC chemokine 2; chemokine CC-1/CC-3; hemofiltrate CC chemokine 1; small-inducible cytokine A14; small inducible cytokine subfamily A (Cys-Cys), member 14; CC-1; CC-3; CKB1; NCC2; SY14; HCC-1; HCC-3; NCC-2; SCYA14; HCC-1(1-74); HCC-1/HCC-3; FLJ16015;

Gene ID

6358

mRNA Refseq

NM_032963

Protein Refseq

NP_116739

MIM

601392

UniProt ID

Q16627

Chromosome Location

17q11.2

Pathway

Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

pathway, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;

Function

chemokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA