

Recombinant Human CASR 293 Cell Lysate

Cat. No. CASR-7825HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for calcium-sensing receptor (CASR) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

CASR calcium-sensing receptor [Homo sapiens]

Official Symbol

CASR

Synonyms

CASR; calcium-sensing receptor; HHC, HHC1, hypocalciuric hypercalcemia 1; extracellular calcium-sensing receptor; FHH; GPRC2A; NSHPT; severe neonatal hyperparathyroidism; parathyroid Ca(2+)-sensing receptor 1; parathyroid cell calcium-sensing receptor; CAR; FIH; HHC; EIG8; HHC1; PCAR1; MGC138441;

Gene ID

846

mRNA Refseq

NM_000388

Protein Refseq

NP_000379

UniProt ID

P41180

Chromosome Location

3q21.1

Pathway

Class C/3 (Metabotropic glutamate/pheromone receptors), organism-specific biosystem; E-cadherin signaling in keratinocytes, organism-specific biosystem; G alpha (i) signalling events, organism-specific biosystem; G alpha (q) signalling events, organism-specific biosystem; GPCR downstream signaling, organism-specific biosystem; GPCR ligand binding, organism-specific biosystem; GPCRs, Class C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Metabotropic glutamate, pheromone, organism-specific biosystem;

Function

G-protein coupled receptor activity; phosphatidylinositol phospholipase C activity; protein binding; receptor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA