

Recombinant Human BCKDHB 293 Cell Lysate

Cat. No. BCKDHB-8493HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for branched chain keto acid dehydrogenase E1, beta polypeptide (BCKDHB), nuclear gene encoding mitochondrial protein, transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

[BCKDHB branched chain keto acid dehydrogenase E1, beta polypeptide \[Homo sapiens \]](#)

Official Symbol

BCKDHB

Synonyms

BCKDHB; branched chain keto acid dehydrogenase E1, beta polypeptide; 2-oxoisovalerate dehydrogenase subunit beta, mitochondrial; maple syrup urine disease; BCKDE1B; BCKDH E1-beta; 2-oxoisovalerate dehydrogenase beta subunit; E1b-beta subunit of the branched-chain complex; branched chain alpha-ketoacid dehydrogenase E1-beta subunit; branched-chain alpha-keto acid dehydrogenase E1 component beta chain; E1B; dJ279A18.1; FLJ17880;

Gene ID

[594](#)

mRNA Refseq

[NM_000056](#)

Protein Refseq

[NP_000047](#)

MIM

[248611](#)

UniProt ID

[P21953](#)

Chromosome Location

6q14.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	2-oxobutanoate degradation I, organism-specific biosystem; 2-oxobutanoate degradation I, conserved biosystem; Branched-chain amino acid catabolism, organism-specific biosystem; Leucine degradation, leucine =>acetoacetate + acetyl-CoA, organism-specific biosystem; Leucine degradation, leucine => acetoacetate + acetyl-CoA, conserved biosystem;
Function	3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring) activity; alpha-ketoacid dehydrogenase activity; carboxy-lyase activity; protein binding; protein complex binding;