

Recombinant Human BCAT1 293 Cell Lysate

Cat. No. BCAT1-8495HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for branched chain aminotransferase 1, cytosolic (BCAT1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [BCAT1 branched chain amino-acid transaminase 1, cytosolic \[Homo sapiens \]](#)

Official Symbol BCAT1

Synonyms BCAT1; branched chain amino-acid transaminase 1, cytosolic; BCT1, branched chain aminotransferase 1, cytosolic; branched-chain-amino-acid aminotransferase, cytosolic; placental protein 18; branched chain aminotransferase 1, cytosolic; BCT1; PP18; BCATC; ECA39; MECA39; PNAS121; DKFZp686E12175;

Gene ID [586](#)

mRNA Refseq [NM_001178091](#)

Protein Refseq [NP_001171562](#)

MIM [113520](#)

UniProt ID [P54687](#)

Chromosome Location 12pter-q12

Pathway Branched-chain amino acid catabolism, organism-specific biosystem; Leucine degradation, leucine =>acetoacetate + acetyl-CoA, organism-specific biosystem; Leucine degradation, leucine => acetoacetate + acetyl-CoA, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem;

Function

L-isoleucine transaminase activity; L-leucine transaminase activity; L-valine transaminase activity; branched-chain-amino-acid transaminase activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA