

Recombinant Human BAG4 293 Cell Lysate

Cat. No. BAG4-8526HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for BCL2-associated athanogene 4 (BAG4) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

BAG4 BCL2-associated athanogene 4 [Homo sapiens]

Official Symbol

BAG4

Synonyms

BAG4; BCL2-associated athanogene 4; BAG family molecular chaperone regulator 4; silencer of death domains; SODD; bcl-2-associated athanogene 4; BAG-family molecular chaperone regulator-4; BAG-4; DKFZp586O2022;

Gene ID

9530

mRNA Refseq

NM_001204878

Protein Refseq

NP_001191807

MIM

603884

UniProt ID

O95429

**Chromosome
Location**

8p11.23

Pathway

Ceramide signaling pathway, organism-specific biosystem; HIV-1 Nef: Negative effector of Fas and TNF-alpha, organism-specific biosystem; TNF receptor signaling pathway, organism-specific biosystem; TNF-alpha/NF-kB Signaling Pathway, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

chaperone binding; receptor signaling protein activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA