

Recombinant Human BAD 293 Cell Lysate

Cat. No. BAD-8529HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for BCL2-associated agonist of cell death (BAD), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

BAD BCL2-associated agonist of cell death [Homo sapiens]

Official Symbol

BAD

Synonyms

BAD; BCL2-associated agonist of cell death; bcl2 antagonist of cell death; BBC2; BCL2L8; bcl2-L-8; BCL2-binding protein; bcl-2-like protein 8; BCL2-binding component 6; bcl-2-binding component 6; BCL-X/BCL-2 binding protein; BCL2-antagonist of cell death protein; bcl-XL/Bcl-2-associated death promoter;

Gene ID

572

mRNA Refseq

NM_004322

Protein Refseq

NP_004313

MIM

603167

UniProt ID

Q92934

Chromosome Location

11q13.1

Pathway

AKT phosphorylates targets in the cytosol, organism-specific biosystem; Activation of BAD and translocation to mitochondria, organism-specific biosystem; Activation of BH3-only proteins, organism-specific biosystem; Acute myeloid leukemia, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; Acute myeloid leukemia, conserved biosystem; Adaptive Immune System, organism-specific biosystem; Alpha-synuclein signaling, organism-specific biosystem;

Function

cysteine-type endopeptidase activator activity involved in apoptotic process; lipid binding; phospholipid binding; protein binding; protein heterodimerization activity; protein kinase binding; protein phosphatase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA