

Recombinant Human BACH1 293 Cell Lysate

Cat. No. BACH1-8532HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for BTB and CNC homology 1, basic leucine zipper transcription factor 1 (BACH1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

BACH1 BTB and CNC homology 1, basic leucine zipper transcription factor 1 [Homo sapiens]

Official Symbol

BACH1

Synonyms

BACH1; BTB and CNC homology 1, basic leucine zipper transcription factor 1; transcription regulator protein BACH1; BACH 1; HA2303; BTB and CNC homolog 1; basic region leucine zipper transcriptional regulator BACH1; BACH-1;

Gene ID

571

mRNA Refseq

NM_206866

Protein Refseq

NP_996749

MIM

602751

UniProt ID

O14867

Chromosome Location

21q22.1

Function

RNA polymerase II core promoter proximal region sequence-specific DNA binding transcription factor activity involved in negative regulation of transcription; RNA polymerase II core promoter proximal region sequence-specific DNA binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

transcription factor activity involved in negative regulation of transcription; heme binding; protein dimerization activity; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity;