

Recombinant Human LIF protein

Cat. No. LIF-22H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human LIF protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	180
Description	Leukemia inhibitory factor (LIF) is a member of Interleukin 6 family. This protein is mainly expressed in the trophectoderm of the developing embryo, with its receptor LIFR expressed throughout the inner cell mass. LIF has the capacity to induce terminal differentiation in leukemic cells. Its activities include the induction of hematopoietic differentiation in normal and myeloid leukemia cells, the induction of neuronal cell differentiation, and the stimulation of acute-phase protein synthesis in hepatocytes. LIF is used in mouse embryonic stem cell culture, because that removal of LIF pushes stem cells toward differentiation, but they retain their proliferative potential or pluripotency. It is also used in phase II clinical trial, which can assist embryo implantation in women who have failed to become pregnant despite assisted reproductive technologies (ART). Mature human LIF (180 a.a.) shares 78 %, 82 %, 91 %, 88 % and 87 % a.a. sequence identity with mouse, rat, canine, bovine, and porcine LIF, respectively.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	dose-dependent proliferation of human TF-1 cells is less than 0.1 ng/ml, corresponding to a specific activity of $> 1.0 \times 10^7$ IU/mg.
Molecular Mass	Approximately 19.7 kDa, a single non-glycosylated polypeptide chain containing 180 amino acids.
AA Sequence	SPLPITPVNATCAIRHPCHNNLMNQIRSQAQLNGSANALFILYYTAQGEPFPNNLDK LCGPNVTDFPPFHANGTEKAKLVELYRIVVYLGTSLGNITRDQKILNPSALSLSKLN ATADILRGLLSNVLCRLCSKYHVGHVDTYGPDTSGKDVFQKKKLGCQLLGKYKQII AVLAQAF
Endotoxin	Less than 1 EU/μg of rHuLIF as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-0.2 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	LIF
Official Symbol	LIF

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	LIF; leukemia inhibitory factor; CDF; cholinergic differentiation factor; DIA; differentiation inhibitory activity; differentiation inducing factor; hepatocyte stimulating factor III; HILDA; human interleukin in DA cells; D factor; melanoma-derived LPL inhibitor; differentiation-inducing factor; hepatocyte-stimulating factor III; differentiation-stimulating factor; MLPLI;
Gene ID	3976
mRNA Refseq	NM_001257135
Protein Refseq	NP_001244064
MIM	159540
UniProt ID	P15018
SDS-PAGE of LIF-22H	

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA