

Recombinant Human AOC2 293 Cell Lysate

Cat. No. AOC2-8823HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for amine oxidase, copper containing 2 (retina-specific) (AOC2), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [AOC2 amine oxidase, copper containing 2 \(retina-specific\) \[Homo sapiens \]](#)

Official Symbol AOC2

Synonyms AOC2; amine oxidase, copper containing 2 (retina-specific); retina-specific copper amine oxidase; DAO2; RAO; SSAO; semicarbazide-sensitive amine oxidase;

Gene ID [314](#)

mRNA Refseq [NM_009590](#)

Protein Refseq [NP_033720](#)

MIM [602268](#)

UniProt ID [O75106](#)

Chromosome Location 17q21

Pathway Glycine, serine and threonine metabolism, organism-specific biosystem; Glycine, serine and threonine metabolism, conserved biosystem; Metabolic pathways, organism-specific biosystem; Phenylalanine metabolism, organism-specific biosystem; Phenylalanine metabolism, conserved biosystem; Tyrosine metabolism, organism-specific biosystem; Tyrosine metabolism, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

aliphatic-amine oxidase activity; aminoacetone:oxygen oxidoreductase(deaminating) activity; copper ion binding; electron carrier activity; metal ion binding; oxidoreductase activity; phenethylamine:oxygen oxidoreductase (deaminating) activity; primary amine oxidase activity; quinone binding; tryptamine:oxygen oxidoreductase (deaminating) activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA