

Recombinant Human ALG2 293 Cell Lysate

Cat. No. ALG2-8906HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for asparagine-linked glycosylation 2, alpha-1,3-mannosyltransferase homolog (*S. cerevisiae*) (ALG2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name ALG2 asparagine-linked glycosylation 2, alpha-1,3-mannosyltransferase homolog (S. cerevisiae) [Homo sapiens]

Official Symbol ALG2

Synonyms ALG2; asparagine-linked glycosylation 2, alpha-1,3-mannosyltransferase homolog (S. cerevisiae); asparagine linked glycosylation 2 homolog (yeast, alpha 1,3 mannosyltransferase); alpha-1,3/1,6-mannosyltransferase ALG2; CDGLi; FLJ14511; hALPG2; NET38; homolog of yeast ALG2; alpha-1,3-mannosyltransferase ALG2; asparagine-linked glycosylation protein 2 homolog; GDP-Man:Man(1)GlcNAc(2)-PP-dolichol mannosyltransferase; GDP-Man:Man(1)GlcNAc(2)-PP-Dol alpha-1,3-mannosyltransferase; GDP-Man:Man(2)GlcNAc(2)-PP-Dol alpha-1,6-mannosyltransferase; asparagine-linked glycosylation 2 homolog (yeast, alpha-1,3-mannosyltransferase); asparagine-linked glycosylation 2 homolog (S. cerevisiae, alpha-1,3-mannosyltransferase);

Gene ID 85365

mRNA Refseq NM_033087

Protein Refseq NP_149078

MIM 607905

UniProt ID Q9H553

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	9q31.1
Pathway	Asparagine N-linked glycosylation, organism-specific biosystem; Biosynthesis of the N-glycan precursor (dolichol lipid-linked oligosaccharide, LLO) and transfer to a nascent protein, organism-specific biosystem; Metabolic pathways, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; N-Glycan biosynthesis, organism-specific biosystem; N-Glycan biosynthesis, conserved biosystem; N-glycan precursor biosynthesis, organism-specific biosystem;
Function	GDP-Man:Man1GlcNAc2-PP-Dol alpha-1,3-mannosyltransferase activity; alpha-1,3-mannosyltransferase activity; calcium-dependent protein binding; calcium-dependent protein binding; protein N-terminus binding; protein anchor; protein binding; protein heterodimerization activity; transferase activity, transferring glycosyl groups;