Recombinant Human AIFM3 293 Cell Lysate

Cat. No. AIFM3-8952HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	HEK293
Description	Antigen standard for apoptosis-inducing factor, mitochondrion-associated, 3 (AIFM3), nuclear gene encoding mitochondrial protein, transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	AIFM3 apoptosis-inducing factor, mitochondrion-associated, 3 [Homo sapiens]
Official Symbol	AIFM3
Synonyms	AIFM3; apoptosis-inducing factor, mitochondrion-associated, 3; apoptosis-inducing factor 3; AIFL; FLJ30473; apoptosis-inducing factor like; FLJ45137;
Gene ID	150209
mRNA Refseq	NM_001018060
Protein Refseq	NP_001018070
UniProt ID	Q96NN9
Chromosome Location	22q11.21
Function	2 iron, 2 sulfur cluster binding; cysteine-type endopeptidase activator activity involved in apoptotic process; flavin adenine dinucleotide binding; metal ion binding; oxidoreductase activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA