

Recombinant Human Mediator Complex Subunit 4, His-tagged

Cat. No. MED4-2406H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human MED4, fused to His-tag at C-terminus, was expressed in E.coli and purified by using conventional chromatography techniques, 30.7kDa (278aa).
Species	Human
Source	E.coli
Description	Mediator complex subunit 4(MED4) is also known as mediator of RNA polymerase II transcription subunit 4 or vitamin D3 receptor-interacting protein complex 36 kDa component (DRIP36). This protein is a component of the vitamin D receptor-interacting protein (DRIP) complex which functions as a nuclear receptor coactivator. The DRIP complex is capable of activating nuclear receptors in a ligand-dependent manner.
Form	Liquid. 20mM Tris-HCl buffer (pH8.0) containing 20% glycerol,1mM DTT, 100mM NaCl.
Purity	> 80% by SDS – PAGE.
Concentration	0.5 mg/ml (determined by Bradford assay).
Sequences Of Amino Acids	MAASSSGEKE KERLGGGLGV AGGNSTRERL LSALEDLEVL SRELIEMLA SRNQKLLQAG EENQVLELLI HRDGEFQELM KLALNQGKIH HEMQVLEKEV EKRDGDIQQL QKQLKEAEQI LATAVYQAKE KLKSIEKARK GAISSEEIIK YAHRIASNA VCAPLTWVPG DPRRPYPTDL EMRSGLLGQM NNPSTNGVNG

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

HLPGDALAAG RLPDVLAPQY PWQSNDSMSN MLPPNHSSDF LLEPPGHNKE
DEDDVEIMST DSSSSSESSED LEHHHHHH.

Storage

Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

GENE INFORMATION
Gene Name

MED4 mediator complex subunit 4 [Homo sapiens]

Synonyms

mediator complex subunit 4; VDRIP; DRIP36; TRAP36; HSPC126; FLJ10956; RP11-90M2.2; MED4; mediator of RNA polymerase II transcription subunit 4; ARC36; p36 TRAP/SMCC/PC2 subunit; TRAP/SMCC/PC2 subunit p36 subunit; vitamin D receptor interacting protein; activator-recruited cofactor 36 kDa component; mediator of RNA polymerase II transcription, subunit 4 homolog; vitamin D3 receptor-interacting protein complex 36 kDa component ;mediator of RNA polymerase II transcription, subunit 4 homolog (S. cerevisiae); Activator-recruited cofactor 36 kDa component; Vitamin D3 receptor-interacting protein complex 36 kDa component

Gene ID

29079

mRNA Refseq

NM_014166

Protein Refseq

NP_054885

MIM

605718

UniProt ID

Q9NPJ6

**Chromosome
Location**

13q14.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Gene Expression
Function	RNA polymerase II transcription mediator activity; ligand-dependent nuclear receptor transcription coactivator activity; protein binding; receptor activity; thyroid hormone receptor binding; transcription activator activity; transcription cofactor activity; vitamin D receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA