

Recombinant Human BCR cell lysate

Cat. No. BCR-166HCL Lot. No. (See product label)

SPECIFICATION

Product Overview

Species	Human
Description	A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene.
Size	100 ul
Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)
Applications	Western Blot;

GENE INFORMATION

Gene Name	BCR breakpoint cluster region [Homo sapiens]
-----------	--

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	BCR
Synonyms	BCR; breakpoint cluster region; BCR1, D22S11; breakpoint cluster region protein; ALL; CML; D22S662; PHL; renal carcinoma antigen NY-REN-26; BCR1; D22S11; FLJ16453;
Gene ID	613
mRNA Refseq	NM_004327
Protein Refseq	NP_004318
MIM	151410
UniProt ID	P11274
Chromosome Location	22q11
Pathway	Chronic myeloid leukemia, organism-specific biosystem; Chronic myeloid leukemia, conserved biosystem; Pathways in cancer, organism-specific biosystem; Regulation of RAC1 activity, organism-specific biosystem; Regulation of RhoA activity, organism-specific biosystem; Rho GTPase cycle, organism-specific biosystem; Signal Transduction, organism-specific biosystem;
Function	ATP binding; GTPase activator activity; Rac GTPase activator activity; Rho guanyl-nucleotide exchange factor activity; guanyl-nucleotide exchange factor activity; kinase activity; nucleotide binding; protein serine/threonine kinase activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA