

Recombinant Human GPD2 cell lysate

Cat. No. GPD2-733HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Description	Mitochondrial glycerophosphate dehydrogenase (EC 1.1.99.5), or GPD2, is located on the outer surface of the inner mitochondrial membrane and catalyzes the unidirectional conversion of glycerol-3-phosphate (G-3-P) to dihydroxyacetone phosphate (DHAP) with concomitant reduction of the enzyme-bound FAD. Together with a cytosolic NAD-linked GPD (GPD1; MIM 138420), GPD2 forms the glycerol phosphate shuttle, which uses the interconversion of G-3-P and DHAP to transfer reducing equivalents into mitochondria, resulting in the reoxidation of NADH formed during glycolysis.
Size	100 ug
Recommended Usage	Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95°C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Storage Buffer	In modified RIPA Lysis Buffer.
Applications	Western Blot;Immunoprecipitation;

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	GPD2 glycerol-3-phosphate dehydrogenase 2 (mitochondrial) [Homo sapiens]
Official Symbol	GPD2
Synonyms	GPD2; glycerol-3-phosphate dehydrogenase 2 (mitochondrial); glycerol-3-phosphate dehydrogenase, mitochondrial; GPD-M; mtGPD; GPDH-M; mitochondrial glycerophosphate dehydrogenase; GDH2; GPDM; mGPDH;
Gene ID	2820
mRNA Refseq	NM_000408
Protein Refseq	NP_000399
MIM	138430
UniProt ID	P43304
Chromosome Location	2q24.1
Pathway	Fatty Acid Beta Oxidation, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Glycerophospholipid metabolism, organism-specific biosystem; Glycerophospholipid metabolism, conserved biosystem; Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem; glycerol-3-phosphate shuttle, organism-specific biosystem;
Function	calcium ion binding; glycerol-3-phosphate dehydrogenase [NAD+] activity; glycerol-3-phosphate dehydrogenase activity; oxidoreductase activity; sn-glycerol-3-phosphate:ubiquinone-8 oxidoreductase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA