

Recombinant Human SIRT2, His-tagged

Cat. No. SIRT2-929H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human SIRT2 protein, fused to His-tag at N-terminus, was expressed in <i>E.coli</i> and purified by using conventional chromatography techniques. MW = 41.7kDa (372aa).
Species	Human
Source	E.coli
ProteinLength	1-352 aa
Description	SIRT2 (NAD-dependent deacetylase sirtuin-2) is a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes and are involved in diverse processes, including transcriptional regulation, cell cycle progression, DNA-damage repair and aging. SIRT2 is a NAD-dependent deacetylase, which deacetylates the "Lys-40" of alpha-tubulin.
Form	Liquid. 20mM Tris-HCl buffer (pH8.0) containing 30% glycerol, 2mM DTT, 200mM NaCl, 0.5mM EDTA.
Purity	> 90% by SDS - PAGE.
Concentration	0.25mg/ml (determined by Bradford assay).
Sequences Of Amino	MGSSHHHHHH SSGLVPRGSH MDFLRNLFSSQ TSLSGSQKER LLELTLEGV ARY

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Acids

MQSERCR RVICLVGAGI STSAGIPDFR SPSTGLYDNL EKYHLPYPEA
 IFEISYFKKH PEPFFALAKE LYPGQFKPTI CHYFMRLKLD KGLLLRCYTQ
 NIDTLERIAG LEQE DLVEAH GTFYTSHCVSASCRHEYPLS WMKEKIFSEV
 TPKCEDCQSL VKPDI VF FGE SLPARFFSCM QSDFLKVDLL LVMGTSLQVQ
 PFASLISKAP LSTPRLLINK EK A GQ SDPFL GMIMGLGGGM DFDSKKAYRD
 VAWLGECDQG CLALAELLGW KKE LEDL VRR EHASIDAQSG AGVNPSTSA
 SPKKSPPPAK DEARTTEREK PQ.

Storage

Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

GENE INFORMATION
Gene Name

SIRT2 sirtuin (silent mating type information regulation 2 homolog) 2 (S. cerevisiae) [Homo sapiens]

Synonyms

sirtuin (silent mating type information regulation 2 homolog) 2 (S. cerevisiae); SIR2; SIR2L; SIR2L2; SIRT2; sirtuin 2; sirtuin type 2; sir2-related protein type 2; silent information regulator 2; silencing information regulator 2-like; EC 3.5.1.-; SIR2-like; SIR2-like protein 2; sirtuin (silent mating type information regulation 2, S.cerevisiae, homolog) 2

Gene ID

22933

mRNA Refseq

NM_012237

Protein Refseq

NP_036369

MIM

604480

UniProt ID

Q8IXJ6

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Chromosome
Location**

19q13

Function

NAD binding; NAD+ ADP-ribosyltransferase activity; NAD-dependent histone deacetylase activity; histone acetyltransferase binding; histone deacetylase binding; hydrolase activity; metal ion binding; transcription factor binding; tubulin deacetylase activity; ubiquitin binding; zinc ion binding

**PDB rendering based
on 1j8f.**

