

Ni-NTA Polar IMAC PMMA resin

Cat. No. IMAC-59P **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Ni-NTA Polar IMAC PMMA resin packing is a metal-affinity packing suitable for purifying recombinant proteins with histidine (His) tags, which are widely used in the biotechnology field. The filler is based on hydrophilic polymethacrylate microspheres with A particle size of 30 µm or 60 µm and a pore size of 800 Å, and has high physical and chemical stability. The surface of the filler has a high degree of hydrophilicity, which avoids non-specific adsorption with biological samples to the greatest extent. On the hydrophilic surface, carboxylic acid groups that can chelate polyvalent metal ions are linked with a proprietary connector.
Form	50% (v/v) suspension, preserved in 20% ethanol
Matrix	Hydrophilic polymethacrylate
Average particle size	30 µm; 60 µm
Ligand	NTA
pH stability range	Base filler Polar MC-CA Excel: 1-13; Metal ion chelating filler Polar MC-Ni Excel: 1-13
Capacity	Polar MC30-CA Excel >120 µeq. carboxylic group /mL Polar MC30-Ni Excel 40-60 µeq.Ni++ /mL Polar MC60-CA Excel >100 µeq. carboxylic group /mL Polar MC60-Ni Excel 40-60 µeq.Ni++ /mL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Histidine labeled protein (Histidine protein) loading	Polar MC30-Ni Excel, ~20 mg/mL Polar MC60-Ni Excel, ~15 mg/mL
Operating Temperature	≤40°C
Operating pressure	≤30 MPa (30 bar)
Mobile phase compatibility	Aqueous solution, or a mixture of water and acetonitrile, acetone, or methanol. Commonly used buffers: phosphate, trimethylol aminomethane (Tris) and acetate, also compatible with 6M guanidine hydrochloride, 8M urea, 0.1 M EDTA, 10 mM DTT
Linear flow rate	≤1800 cm/h
Storage	Store at 20% ethanol, 4-25°C.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA