

Rabbit Anti-Human IgG Fc Secondary Polyclonal Antibody, HRP

Cat. No. CPB-054RH **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source Rabbit

Immunogen Description

Antibodies, also known as Immunoglobulins (Ig), are gamma globulin proteins that are found in blood or other bodily fluids of vertebrates, and are used by the immune system to identify and neutralize foreign objects, such as bacteria and viruses. Antibodies are heavy globular plasma proteins. They have sugar chains added to some of their amino acid residues. In other words, antibodies are glycoproteins. Though the general structure of all antibodies is very similar, a small region at the tip of the protein is extremely variable, allowing millions of antibodies with slightly different tip structures, or antigen binding sites, to exist. This region is known as the hypervariable region. Each of these variants can bind to a different target, known as an antigen. Immunoglobulin G (IgG) is a monomeric immunoglobulin, built of two heavy chains and two light chains. Each IgG has two antigen binding sites. It is the most abundant immunoglobulin and is approximately equally distributed in blood and in tissue liquids, constituting 75% of serum immunoglobulins in humans. IgG molecules are synthesized and secreted by plasma B cells. Antibodies can come in different varieties known as isotypes or classes. In placental mammals there are five antibody isotypes known as IgA, IgD, IgE, IgG and IgM. They are each named with an "Ig" prefix that stands for immunoglobulin, another name for antibody, and differ in their biological properties, functional locations and ability to deal with different

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

antigens. The basic functional unit of each antibody is an immunoglobulin (Ig) monomer (containing only one Ig unit); secreted antibodies can also be dimeric with two Ig units as with IgA, tetrameric with four Ig units like teleost fish IgM, or pentameric with five Ig units, like mammalian IgM. The variable parts of an antibody are its V regions, and the constant part is its C region.

Immunogen	Human IgG.
Antibody Type	HRP conjugated Rabbit Polyclonal Antibody.
Ig Type	Rabbit IgG.
Formulation	0.2 µm filtered solution in PBS with 5% trehalose.
Preparation	Produced in rabbits immunized with purified human IgG. This polyclonal antibody was purified by antigen affinity chromatography and conjugated with horseradish-peroxidase (HRP).
Specificity	Human IgG Fc. Has cross-reactivity with goat IgG in ELISA. No cross-reactivity with human IgM , rabbit IgG and mouse IgG in ELISA.
Western Blot	This antibody can be used at 0.05 - 0.2 µg/mL in WB. Optimal dilutions should be determined by each laboratory for each application.
DirectELISA	This antibody can be used at 0.05 - 0.2 µg/mL in ELISA. Optimal dilutions should be determined by each laboratory for each application.
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -70°C. Preservative-Free. Sodium azide is recommended

to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA