

Active Human CXCL9 Reference Standard

Cat. No. CXCL9-10H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Form	Liquid
Bio-activity	The level of CXCL9 in the serum was quantified using ELISA.
Usage	For research use only, not for diagnostic or therapeutic use.
Quality Control Test	Generated from a human serum pool. This lot has been tested by standard laboratory procedures and found to be negative for HIV 1, HIV 2, Hepatitis B and Hepatitis C.
Storage	Stored at -20 °C, 4 °C for less than seven days, frozen for greater than seven days.
Concentration	3.97 ng/ml
Preservative	0.1% sodium azide

GENE INFORMATION

Gene Name	CXCL9 chemokine (C-X-C motif) ligand 9 [Homo sapiens]
Official Symbol	CXCL9
Synonyms	CXCL9; chemokine (C-X-C motif) ligand 9; CMK, MIG, monokine induced by gamma

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	interferon; C-X-C motif chemokine 9; crg 10; Humig; SCYB9; small-inducible cytokine B9; gamma-interferon-induced monokine; monokine induced by gamma interferon; monokine induced by interferon-gamma; CMK; MIG; crg-10;
Gene ID	4283
mRNA Refseq	NM_002416
Protein Refseq	NP_002407
MIM	601704
UniProt ID	Q07325
Chromosome Location	4q21
Pathway	CXCR3-mediated signaling events, organism-specific biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
Function	chemokine activity; cytokine activity;