

Active Recombinant Human IL2, His tagged

Cat. No. IL2-501H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human Interleukin-2/IL-2 is produced by our mammalian expression system in human cells. The target protein is expressed with sequence (Pro22-Thr153) of Human IL-2 fused with a polyhistidine tag at the C-terminus.
Species	Human
Source	Human Cells
ProteinLength	22-153 a.a.
Description	Recombinant Human Interleukin-2 is a highly purified protein with a molecular weight of approximately 15,300 Daltons. The chemical name is des-alanyl-1, serine-125 Human Interleukin-2. It is produced by recombinant DNA technology using a genetically engineered E. coli strain containing an analog of the human interleukin-2 gene. Genetic engineering techniques were used to modify the Human IL-2 gene, and the resulting expression clone encodes a modified Human IL-2. This recombinant form differs from native Interleukin-2 in following ways: 1) it is not glycosylated; 2) the molecule has no N-terminal alanine; 3) the molecule has serine substituted for cysteine at amino acid position 125; 4) the aggregation state of molecule is likely to be different from that of native IL-2.
Form	Lyophilized from a 0.2 µM filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Bio-activity	>1.0 x 10 ⁷ U/mg

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 0.1 EU/μg.
Purity	Greater than 95% as determined by SEC-HPLC and reducing SDS-PAGE.
Storage	Lyophilized protein should be stored at< -20°C, though stable at room temperature for 3>Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at< -20°C for 5>
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/ml. Dissolve the lyophilized protein in 1X PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

GENE INFORMATION

Gene Name	IL2 interleukin 2 [Homo sapiens]
Official Symbol	IL2
Synonyms	IL2; interleukin 2; interleukin-2; IL 2; T cell growth factor; TCGF; aldesleukin; involved in regulation of T-cell clonal expansion; IL-2; lymphokine;
Gene ID	3558
mRNA Refseq	NM_000586
Protein Refseq	NP_000577
MIM	147680
UniProt ID	P60568

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	4q26-q27
Pathway	Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Calcium signaling in the CD4+ TCR pathway, organism-specific biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem;
Function	carbohydrate binding; cytokine activity; glycosphingolipid binding; growth factor activity; interleukin-2 receptor binding; interleukin-2 receptor binding; kappa-type opioid receptor binding; kinase activator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA