

Human Tumour Necrosis Factor alpha (rDNA derived) Reference standard

Cat. No. TNF-43H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

The Human TNF Reference Standard was for human Tumour Necrosis Factor-alpha (TNF- α) following evaluation in an International Collaborative Study. This preparation is intended for use as the primary biological reference standard in bioassays of human TNF- α .

Species

Human

Enzyme Unit

The preparation has been assigned a potency of 43,000 International Units of biological activity per ampoule.

Usage

No attempt should be made to weigh out any portion of the freeze-dried material prior to reconstitution. Dissolve the total contents of the ampoule in 1.0 ml of sterile distilled water. This solution will contain TNF- α at a concentration of 43,000 International Units/ml. Use carrier protein where extensive dilution is required.

Notes

This preparation is not for administration to humans. The preparation contains material of human origin, and either the final product or the source materials, from which it is derived, have been tested and found negative for HBsAg, anti-HIV and HCV RNA. As with all materials of biological origin, this preparation should be regarded as potentially hazardous to health. It should be used and discarded according to your own laboratory's safety procedures. Such safety procedures should include the wearing of protective gloves and avoiding the generation of aerosols. Care should be exercised in opening ampoules or vials, to avoid cuts.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Stability

Reference materials are held at Creative Biomart within assured, temperature-controlled storage facilities and they should be stored on receipt as indicated on the label. Accelerated degradation studies have indicated that this material is suitably stable, when stored at -20°C or below, for the assigned values to remain valid until the material is withdrawn or replaced. These studies have also shown that the material is suitably stable for shipment at ambient temperature without any effect on the assigned values. Once reconstituted, diluted or aliquoted, users should determine the stability of the material according to their own method of preparation, storage and use. Users who have data supporting any deterioration in the characteristics of any reference preparation are encouraged to contact Creative Biomart.

Storage

For economy of use, it is recommended that the solution be sub divided into several small aliquots and stored at -40°C or below. Avoid repeated thawing/freezing. Unopened ampoules should be stored at -20°C. Please note: because of the inherent stability of lyophilized material, Creative Biomart may ship these materials at ambient temperature.

GENE INFORMATION

Gene Name

TNF tumor necrosis factor [Homo sapiens];

Official Symbol

TNF

Synonyms

TNF; tumor necrosis factor; TNFA, tumor necrosis factor (TNF superfamily, member 2); DIF; TNF superfamily; member 2; TNF alpha; TNFSF2; TNF-a; cachectin; APC1 protein; TNF, monocyte-derived; TNF, macrophage-derived; TNF superfamily, member 2; tumor necrosis factor alpha; tumor necrosis factor-alpha; tumor necrosis factor ligand superfamily member 2; TNFA; TNF-alpha;

Gene ID

7124

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_000594
Protein Refseq	NP_000585
MIM	191160
UniProt ID	P01375
Chromosome Location	6p21.3
Pathway	Adipocytokine signaling pathway, organism-specific biosystem; Adipocytokine signaling pathway, conserved biosystem; Adipogenesis, organism-specific biosystem; African trypanosomiasis, organism-specific biosystem; African trypanosomiasis, conserved biosystem; Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem;
Function	cytokine activity; identical protein binding; protease binding; protein binding; transcription regulatory region DNA binding; tumor necrosis factor receptor binding;