

Recombinant Human Basic Fibroblast Growth Factor (rDNA derived) Reference standard

Cat. No. FGF2-105H Lot. No. (See product label)

SPECIFICATION

Product Overview	The bFGF used in this preparation is the human form of the molecule, synthesized in E. coli by recombinant DNA technology.
Species	Human
Source	E.coli
Enzyme Unit	The assigned potency of the bFGF is 1600 International Units (IU) per ampoule.
Usage	No attempt should be made to weigh out any portion of the freeze-dried material prior to reconstitution. The Reference Standard is intended for calibration of local standards. For all practical purposes, each ampoule contains the same quantity of bFGF. The entire contents of each ampoule should be completely dissolved in a known volume of suitable solvent. It is recommended that, when possible, buffer containing carrier protein should be used to minimize loss by surface adsorption. No attempt should be made to weigh out any portion of the freeze-dried material.
Notes	This preparation is not for administration to humans. The material is not of human or bovine origin. As with all materials of biological origin, this preparation should be regarded as potentially hazardous to health. It should be used and discarded according to your own laboratorys safety procedures. Such safety procedures should include the wearing of protective gloves and avoiding the generation of aerosols. Care should be exercised in opening ampoules or vials, to avoid cuts.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Stability

Reference materials are held at Creative Biomart within assured, temperature-controlled storage facilities and they should be stored on receipt as indicated on the label. Accelerated degradation studies have indicated that this material is suitably stable, when stored at -20°C or below, for the assigned values to remain valid until the material is withdrawn or replaced. These studies have also shown that the material is suitably stable for shipment at ambient temperature without any effect on the assigned values. Users who have data supporting any deterioration in the characteristics of any reference preparation are encouraged to contact Creative Biomart.

Storage

The ampoules are shipped at ambient temperature. Unopened ampoules should be stored at -20 °C in the dark. For economy of use, it is recommended that the reconstituted solution be subdivided into several small containers and stored at, or below, -40 °C . Repeated freezing and thawing should be avoided. The ampoules do not contain bacteriostat and solutions of the ampouled material should not be assumed to be sterile. Please note: because of the inherent stability of lyophilized material, Creative Biomart may ship these materials at ambient temperature.

GENE INFORMATION

Gene Name

FGF2 fibroblast growth factor 2 (basic) [Homo sapiens];

Official Symbol

FGF2

Synonyms

FGF2; fibroblast growth factor 2 (basic); FGFB; fibroblast growth factor 2; prostatropin; heparin-binding growth factor 2; basic fibroblast growth factor bFGF; BFGF; FGF-2; HBGF-2;

Gene ID

2247

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_002006
Protein Refseq	NP_001997
MIM	134920
UniProt ID	P09038
Chromosome Location	4q26
Pathway	Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; Downstream signaling of activated FGFR, organism-specific biosystem; Endochondral Ossification, organism-specific biosystem; FGFR ligand binding and activation, organism-specific biosystem; FGFR1 ligand binding and activation, organism-specific biosystem; FGFR1b ligand binding and activation, organism-specific biosystem; FGFR1c ligand binding and activation, organism-specific biosystem;
Function	chemoattractant activity; cytokine activity; fibroblast growth factor binding; fibroblast growth factor receptor binding; growth factor activity; heparin binding; ligand-dependent nuclear receptor transcription coactivator activity; protein binding; protein tyrosine kinase activity; voltage-gated calcium channel activity;