

Human Rantes (rDNA derived) Reference standard

Cat. No. CCL5-120H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Enzyme Unit The assigned potency is 100,000 arbitrary units per ampoule.

Usage No attempt should be made to weigh out any portion of the freeze-dried material prior to reconstitution. Dissolve the total contents of the ampoule in 0.5ml of sterile distilled water. Rinse the ampoule with a further 0.4ml of sterile distilled water and make up the total volume to 1.0ml. This solution will contain RANTES at a concentration of 100,000 units per ml. It is advisable to include a carrier protein during initial reconstitution to prevent loss of material if this does not interfere in the intended assay method.

Notes This preparation is not for administration to humans. The material is not of human or bovine origin. As with all materials of biological origin, this preparation should be regarded as potentially hazardous to health. It should be used and discarded according to your own laboratorys safety procedures. Such safety procedures should include the wearing of protective gloves and avoiding the generation of aerosols. Care should be exercised in opening ampoules or vials, to avoid cuts.

Stability Reference materials are held at Creative Biomart within assured, temperature-controlled storage facilities and they should be stored on receipt as indicated on the label. An arbitrary unitage was assigned to this preparation at the time of production without confirmation of long term stability. Evidence from similar materials prepared

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

by an equivalent process indicates that long term stability is likely to be maintained and the material is suitably stable for shipment at ambient temperature without any effect on the assigned values. Once reconstituted, diluted or aliquoted, users should determine the stability of the material according to their own method of preparation, storage and use. Users who have data supporting any deterioration in the characteristics of any reference preparation are encouraged to contact Creative Biomart.

Storage

For economy of use, it is recommended that the solution be sub divided into aliquots and stored at +4°C for up to 1 week. Unopened ampoules should be stored at –20°C. Please note: because of the inherent stability of lyophilized material, Creative Biomart may ship these materials at ambient temperature.

GENE INFORMATION

Gene Name CCL5 chemokine (C-C motif) ligand 5 [Homo sapiens];

Official Symbol CCL5

Synonyms

CCL5; chemokine (C-C motif) ligand 5; D17S136E, SCYA5, small inducible cytokine A5 (RANTES); C-C motif chemokine 5; beta chemokine RANTES; MGC17164; RANTES; regulated upon activation; normally T expressed; and presumably secreted; SIS delta; SISd; small inducible cytokine subfamily A (Cys Cys); member 5; T cell specific protein p288; T cell specific RANTES protein; TCP228; eoCP; SIS-delta; beta-chemokine RANTES; small-inducible cytokine A5; T-cell specific protein p288; t cell-specific protein P228; T-cell-specific protein RANTES; eosinophil chemotactic cytokine; small inducible cytokine A5 (RANTES); small inducible cytokine subfamily A (Cys-Cys), member 5; regulated upon activation, normally T-expressed, and presumably secreted; SCYA5; D17S136E;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	6352
mRNA Refseq	NM_002985
Protein Refseq	NP_002976
MIM	187011
UniProt ID	P13501
Chromosome Location	17q11.2-q12
Pathway	Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem;
Function	CCR1 chemokine receptor binding; CCR1 chemokine receptor binding; CCR1 chemokine receptor binding; CCR4 chemokine receptor binding; CCR5 chemokine receptor binding; chemoattractant activity; chemokine activity; chemokine activity; chemokine receptor antagonist activity; chemokine receptor binding; heparin binding; phosphatidylinositol phospholipase C activity; phospholipase activator activity; protein homodimerization activity; protein kinase activity; protein self-association; receptor signaling protein tyrosine kinase activator activity;